

# Weighted Hermite-Hadamard-Type Inequalities by Identities Related to Generalizations of Steffensen's Inequality

---

Pečarić, Josip; Perušić Pribanić, Anamarija; Smoljak Kalamir, Ksenija

Source / Izvornik: **Mathematics, 2022, 10**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.3390/math10091505>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:157:855657>

Rights / Prava: [Attribution 4.0 International](#) / [Imenovanje 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2024-06-30**



image not found or type unknown

Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Civil Engineering - FCERI Repository](#)




image not found or type unknown

Article

# Weighted Hermite–Hadamard-Type Inequalities by Identities Related to Generalizations of Steffensen’s Inequality

Josip Pečarić<sup>1</sup>, Anamarija Perušić Pribanić<sup>2,\*</sup>  and Ksenija Smoljak Kalamir<sup>3</sup> 

<sup>1</sup> Croatian Academy of Sciences and Arts, Trg Nikole Šubića Zrinskog 11, 10000 Zagreb, Croatia; pecaric@element.hr

<sup>2</sup> Faculty of Civil Engineering, University of Rijeka, Radmile Matejčić 3, 51000 Rijeka, Croatia

<sup>3</sup> Faculty of Textile Technology, University of Zagreb, Prilaz Baruna Filipovića 28a, 10000 Zagreb, Croatia; ksmoljak@ttf.hr

\* Correspondence: anamarija.perusic@uniri.hr

**Abstract:** In this paper, we obtain some new weighted Hermite–Hadamard-type inequalities for  $(n + 2)$ -convex functions by utilizing generalizations of Steffensen’s inequality via Taylor’s formula.

**Keywords:** weighted Hermite–Hadamard inequality; Steffensen’s inequality; Taylor’s formula;  $n$ -convex functions

**MSC:** 26D15; 26A51



**Citation:** Pečarić, J.; Perušić Pribanić, A.; Smoljak Kalamir, K. Weighted Hermite–Hadamard-Type Inequalities by Identities Related to Generalizations of Steffensen’s Inequality. *Mathematics* **2022**, *10*, 1505. <https://doi.org/10.3390/math10091505>

Academic Editor: Jaan Janno

Received: 30 March 2022

Accepted: 29 April 2022

Published: 1 May 2022

**Publisher’s Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The Hermite–Hadamard inequality is one of the most important mathematical inequalities. It was discovered independently first by Hermite [1] and later by Hadamard [2]. The classical Hermite–Hadamard inequality provides an estimate from below and above the mean value of convex function  $f: [a, b] \rightarrow \mathbb{R}$ . More precisely, we have the following.

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}.$$

To illustrate the importance of the Hermite–Hadamard inequality, let us mention that the Hermite–Hadamard inequality can be considered as the necessary and sufficient condition for convexity of a function. Furthermore, the Hermite–Hadamard inequality has an important role in numerical analysis, mathematical analysis and functional analysis. Various generalizations, extensions and applications of the Hermite–Hadamard inequality have appeared in the literature (see [3–8]).

In this paper, we consider the weighted Hermite–Hadamard inequality for convex functions given in following theorem (see [8–10]).

**Theorem 1.** Let  $p: [a, b] \rightarrow \mathbb{R}$  be a non-negative function. If  $f: [a, b] \rightarrow \mathbb{R}$  is a convex function, then we have the following:

$$f(m) \leq \frac{1}{P(b)} \int_a^b p(x) f(x) dx \leq \frac{b-m}{b-a} f(a) + \frac{m-a}{b-a} f(b)$$

or

$$P(b)f(m) \leq \int_a^b p(x) f(x) dx \leq P(b) \left[ \frac{b-m}{b-a} f(a) + \frac{m-a}{b-a} f(b) \right], \quad (1)$$

where the following is the case.

$$P(t) = \int_a^t p(x) dx \quad \text{and} \quad m = \frac{1}{P(b)} \int_a^b p(x) x dx.$$

In 1918, Steffensen proved the following inequality (see [11]).

**Theorem 2 ([11]).** *Suppose that  $f$  is non-increasing and  $g$  is integrable on  $[a, b]$  with  $0 \leq g \leq 1$  and  $\lambda = \int_a^b g(t)dt$ . Then, we have the following.*

$$\int_{b-\lambda}^b f(t)dt \leq \int_a^b f(t)g(t)dt \leq \int_a^{a+\lambda} f(t)dt. \tag{2}$$

The inequalities are reversed for  $f$  non-decreasing.

Many papers have been devoted to generalizations and refinements of Steffensen’s inequality and its connection to other well-known inequalities such as Gauss–Steffensen’s, Hölder’s, Jensen–Steffensen’s and other inequalities. A complete overview of the results related to Steffensen’s inequality can be found in monographs [12,13].

By using the Mitrinović [14] result in which the inequalities in (2) follow from identities:

$$\begin{aligned} \int_a^{a+\lambda} f(t)dt - \int_a^b f(t)g(t)dt \\ = \int_a^{a+\lambda} [f(t) - f(a + \lambda)][1 - g(t)]dt + \int_{a+\lambda}^b [f(a + \lambda) - f(t)]g(t)dt \end{aligned}$$

and

$$\begin{aligned} \int_a^b f(t)g(t)dt - \int_{b-\lambda}^b f(t)dt \\ = \int_a^{b-\lambda} [f(t) - f(b - \lambda)]g(t)dt + \int_{b-\lambda}^b [f(b - \lambda) - f(t)][1 - g(t)]dt \end{aligned}$$

and using Taylor’s formulae in points  $a$  and  $b$

$$\begin{aligned} f(x) &= \sum_{i=0}^{n-1} \frac{f^{(i)}(a)}{i!} (x - a)^i + \frac{1}{(n - 1)!} \int_a^x f^{(n)}(t)(x - t)^{n-1} dt \\ f(x) &= \sum_{i=0}^{n-1} \frac{f^{(i)}(b)}{i!} (x - b)^i - \frac{1}{(n - 1)!} \int_x^b f^{(n)}(t)(x - t)^{n-1} dt \end{aligned}$$

in paper [15], the authors proved the following identities related to generalizations of Steffensen’s inequality.

**Theorem 3 ([15]).** *Let  $f: [a, b] \rightarrow \mathbb{R}$  be such that  $f^{(n-1)}$  is absolutely continuous for some  $n \geq 2$  and let  $g: [a, b] \rightarrow \mathbb{R}$  be an integrable function such that  $0 \leq g \leq 1$ . Let  $\lambda = \int_a^b g(t)dt$  and let the function  $G_1$  be defined by the following.*

$$G_1(x) = \begin{cases} \int_a^x (1 - g(t))dt, & x \in [a, a + \lambda], \\ \int_x^b g(t)dt, & x \in [a + \lambda, b]. \end{cases}$$

Then, we have the following:

$$\begin{aligned} \int_a^{a+\lambda} f(t)dt - \int_a^b f(t)g(t)dt + \sum_{i=0}^{n-2} \frac{f^{(i+1)}(a)}{i!} \int_a^b G_1(x)(x - a)^i dx \\ = -\frac{1}{(n - 2)!} \int_a^b \left( \int_t^b G_1(x)(x - t)^{n-2} dx \right) f^{(n)}(t)dt \end{aligned} \tag{3}$$

and the following is obtained.

$$\int_a^{a+\lambda} f(t)dt - \int_a^b f(t)g(t)dt + \sum_{i=0}^{n-2} \frac{f^{(i+1)}(b)}{i!} \int_a^b G_1(x)(x-b)^i dx = \frac{1}{(n-2)!} \int_a^b \left( \int_a^t G_1(x)(x-t)^{n-2} dx \right) f^{(n)}(t)dt. \tag{4}$$

**Theorem 4** ([15]). Let  $f: [a, b] \rightarrow \mathbb{R}$  be such that  $f^{(n-1)}$  is absolutely continuous for some  $n \geq 2$  and let  $g: [a, b] \rightarrow \mathbb{R}$  be an integrable function such that  $0 \leq g \leq 1$ . Let  $\lambda = \int_a^b g(t)dt$  and let the function  $G_2$  be defined by the following.

$$G_2(x) = \begin{cases} \int_a^x g(t)dt, & x \in [a, b - \lambda], \\ \int_x^b (1 - g(t))dt, & x \in [b - \lambda, b]. \end{cases}$$

Then, we have the following:

$$\int_a^b f(t)g(t)dt - \int_{b-\lambda}^b f(t)dt + \sum_{i=0}^{n-2} \frac{f^{(i+1)}(a)}{i!} \int_a^b G_2(x)(x-a)^i dx = -\frac{1}{(n-2)!} \int_a^b \left( \int_t^b G_2(x)(x-t)^{n-2} dx \right) f^{(n)}(t)dt \tag{5}$$

and the following is obtained.

$$\int_a^b f(t)g(t)dt - \int_{b-\lambda}^b f(t)dt + \sum_{i=0}^{n-2} \frac{f^{(i+1)}(b)}{i!} \int_a^b G_2(x)(x-b)^i dx = \frac{1}{(n-2)!} \int_a^b \left( \int_a^t G_2(x)(x-t)^{n-2} dx \right) f^{(n)}(t)dt. \tag{6}$$

Since, in this paper, we will deal with  $n$ -convex functions, let us recall the definition of the  $n$ -convex function. For more details on convex functions, we refer the interested reader to [6,8].

Let  $f$  be a real-valued function defined on the segment  $[a, b]$ . The *divided difference* of order  $n$  of the function  $f$  at distinct points  $x_0, \dots, x_n \in [a, b]$  is defined recursively (see [8]) by the following.

$$f[x_i] = f(x_i), \quad (i = 0, \dots, n)$$

$$f[x_0, \dots, x_n] = \frac{f[x_1, \dots, x_n] - f[x_0, \dots, x_{n-1}]}{x_n - x_0}.$$

The value  $f[x_0, \dots, x_n]$  is independent of the order of the points  $x_0, \dots, x_n$ . The definition may be extended to include the case in which some (or all) of the points coincide. Assuming that  $f^{(j-1)}(x)$  exists, we define the following.

$$f[\underbrace{x, \dots, x}_{j\text{-times}}] = \frac{f^{(j-1)}(x)}{(j-1)!}.$$

**Definition 1** ([8]). A function  $f : [a, b] \rightarrow \mathbb{R}$  is said to be  $n$ -convex on  $[a, b]$ ,  $n \geq 0$ , if for all choices of  $(n + 1)$  distinct points in  $[a, b]$ , the  $n$ -th order divided difference of  $f$  satisfies the following.

$$f[x_0, \dots, x_n] \geq 0.$$

Note that 1-convex functions are non-decreasing functions and 2-convex functions are convex functions. An  $n$ -convex function need not to be  $n$ -times differentiable; how-

ever, if  $f^{(n)}$  exists, then  $f$  is  $n$ -convex if and only if  $f^{(n)} \geq 0$ . The following property also holds: if  $f$  is an  $(n + 2)$ -convex function, then there exists the  $n$ -th derivative  $f^{(n)}$ , which is a convex function.

The aim of this paper is to use identities related to generalizations of Steffensen’s inequality, obtained by using Taylor’s formula, to prove new weighted Hermite–Hadamard-type inequalities for  $(n + 2)$ -convex functions.

### 2. Main Results

In this section, applying identities given in Theorems 3 and 4 and the properties of  $n$ -convex functions, we derive new weighted Hermite–Hadamard-type inequalities.

**Theorem 5.** Let  $f: [a, b] \rightarrow \mathbb{R}$  be  $(n + 2)$ -convex on  $[a, b]$  and  $f^{(n-1)}$  absolutely continuous for  $n \geq 2$ . Let  $g: [a, b] \rightarrow \mathbb{R}$  be an integrable function such that  $0 \leq g \leq 1$  and  $\lambda = \int_a^b g(t)dt$ . Let function  $G_1$  be defined by the following.

$$G_1(x) = \begin{cases} \int_a^x (1 - g(t))dt, & x \in [a, a + \lambda], \\ \int_x^b g(t)dt, & x \in [a + \lambda, b]. \end{cases} \tag{7}$$

Then, we have the following:

$$\begin{aligned} P_1(b) \cdot f^{(n)}(m_1) &\leq \\ (n - 2)! \left[ \int_a^b f(t)g(t)dt - \int_a^{a+\lambda} f(t)dt - \sum_{i=0}^{n-2} \frac{f^{(i+1)}(a)}{i!} \int_a^b G_1(x)(x - a)^i dx \right] &\tag{8} \\ &\leq P_1(b) \cdot \left[ \frac{b - m_1}{b - a} f^{(n)}(a) + \frac{m_1 - a}{b - a} f^{(n)}(b) \right], \end{aligned}$$

where the following is the case:

$$P_1(b) = \frac{1}{(n - 1) \cdot n} \left( \int_a^b g(x)(x - a)^n dx - \frac{\lambda^{n+1}}{n + 1} \right) \tag{9}$$

and the following is obtained.

$$m_1 = a + \frac{1}{(n - 1) \cdot n \cdot (n + 1) \cdot P_1(b)} \left( \int_a^b g(x)(x - a)^{n+1} dx - \frac{\lambda^{n+2}}{n + 2} \right). \tag{10}$$

**Proof.** Since  $f^{(n-1)}$  is absolutely continuous, function  $f$  satisfies the conditions of Theorem 3. Therefore, identity (3) holds.

From condition  $0 \leq g \leq 1$ , function  $G_1$  defined by (7) is non-negative. Hence, for every  $n \geq 2$ , we have the following.

$$\int_t^b G_1(x)(x - t)^{n-2} dx \geq 0, \quad t \in [a, b].$$

Define

$$p(t) = \int_t^b G_1(x)(x - t)^{n-2} dx.$$

Since the function  $f$  is  $(n + 2)$ -convex, function  $f^{(n)}$  is convex. Furthermore, function  $p$  is non-negative, so we can apply Theorem 1 and obtain the following inequality:

$$\begin{aligned} P_1(b) \cdot f^{(n)}(m_1) &\leq \int_a^b \left( \int_t^b G_1(x)(x - t)^{n-2} dx \right) f^{(n)}(t) dt \\ &\leq P_1(b) \cdot \left[ \frac{b - m_1}{b - a} f^{(n)}(a) + \frac{m_1 - a}{b - a} f^{(n)}(b) \right], \end{aligned} \tag{11}$$

where  $P_1(b)$  and  $m_1$  are given by

$$P_1(b) = \int_a^b \left( \int_t^b G_1(x)(x-t)^{n-2} dx \right) dt$$

and

$$m_1 = \frac{1}{P_1(b)} \int_a^b \left( \int_t^b G_1(x)(x-t)^{n-2} dx \right) t dt.$$

By calculating  $P_1(b)$  and  $m_1$ , we obtain the following:

$$\begin{aligned} P_1(b) &= \int_a^b \left( \int_t^b G_1(x)(x-t)^{n-2} dx \right) dt \\ &= \int_a^{a+\lambda} \left( \int_a^x (1-g(s)) ds \right) \frac{(x-a)^{n-1}}{n-1} dx + \int_{a+\lambda}^b \left( \int_x^b g(s) ds \right) \frac{(x-a)^{n-1}}{n-1} dx \\ &= \int_a^{a+\lambda} \frac{(x-a)^n}{n-1} dx + \lambda \cdot \int_{a+\lambda}^b \frac{(x-a)^{n-1}}{n-1} dx - \int_a^b \left( \int_a^x g(s) ds \right) \frac{(x-a)^{n-1}}{n-1} dx \\ &= \frac{-\lambda^{n+1}}{(n-1) \cdot n \cdot (n+1)} + \int_a^b g(x) \frac{(x-a)^n}{(n-1) \cdot n} dx \end{aligned}$$

and

$$\begin{aligned} m_1 &= \frac{1}{P_1(b)} \int_a^b \left( \int_t^b G_1(x)(x-t)^{n-2} dx \right) t dt \\ &= \frac{1}{P_1(b)} \int_a^b G_1(x) \left( \int_a^x (x-t)^{n-2} \cdot t dt \right) dx \\ &= \frac{1}{P_1(b)} \int_a^b G_1(x) \left( t \cdot \frac{-(x-t)^{n-1}}{n-1} \Big|_a^x + \int_a^x \frac{(x-t)^{n-1}}{n-1} dt \right) dx \\ &= \frac{1}{P_1(b)} \int_a^b G_1(x) \left( \frac{a \cdot (x-a)^{n-1}}{n-1} + \frac{(x-a)^n}{(n-1) \cdot n} \right) dx \\ &= a + \frac{1}{P_1(b)} \int_a^b G_1(x) \frac{(x-a)^n}{(n-1) \cdot n} dx \\ &= a + \frac{1}{P_1(b)} \left( \frac{-\lambda^{n+2}}{(n-1) \cdot n \cdot (n+1) \cdot (n+2)} + \int_a^b g(x) \frac{(x-a)^{n+1}}{(n-1) \cdot n \cdot (n+1)} dx \right). \end{aligned}$$

Using identity (3) for the middle part of the inequality (11), inequality (11) becomes inequality (8). Hence, the proof is completed.  $\square$

**Theorem 6.** Let  $f: [a, b] \rightarrow \mathbb{R}$  be  $(n+2)$ -convex on  $[a, b]$  and  $f^{(n-1)}$  absolutely continuous for  $n \geq 2$ . Let  $g: [a, b] \rightarrow \mathbb{R}$  be an integrable function such that  $0 \leq g \leq 1$  and  $\lambda = \int_a^b g(t) dt$ . Let function  $G_1$  be defined by (7). If the following is the case:

$$\int_a^t G_1(x)(x-t)^{n-2} dx \leq 0, \quad t \in [a, b],$$

then we have the following:

$$\begin{aligned} &P_2(b) \cdot f^{(n)}(m_2) \leq \\ &(n-2)! \left[ \int_a^b f(t)g(t)dt - \int_a^{a+\lambda} f(t)dt - \sum_{i=0}^{n-2} \frac{f^{(i+1)}(b)}{i!} \int_a^b G_1(x)(x-b)^i dx \right] \tag{12} \\ &\leq P_2(b) \cdot \left[ \frac{b-m_2}{b-a} f^{(n)}(a) + \frac{m_2-a}{b-a} f^{(n)}(b) \right], \end{aligned}$$

where

$$P_2(b) = \frac{1}{(n-1) \cdot n} \left( \frac{(a-b)^{n+1} - (a+\lambda-b)^{n+1}}{n+1} + \int_a^b g(x)(x-b)^n dx \right)$$

and

$$m_2 = b + \frac{1}{(n-1) \cdot n \cdot (n+1) \cdot P_2(b)} \times \left( \frac{(a-b)^{n+2} - (a+\lambda-b)^{n+2}}{n+2} + \int_a^b g(x)(x-b)^{n+1} dx \right).$$

**Proof.** If we assume the following:

$$\int_a^t G_1(x)(x-t)^{n-2} dx \leq 0, \quad t \in [a, b]$$

then we have the following.

$$-\int_a^t G_1(x)(x-t)^{n-2} dx \geq 0, \quad t \in [a, b].$$

Now similarly to the proof of Theorem 5 using the following non-negative function:

$$p(t) = -\int_a^t G_1(x)(x-t)^{n-2} dx$$

and identity (4), we obtain inequality (12). Similarly, we calculate the expressions for  $P_2(b)$  and  $m_2$  and obtain the following:

$$\begin{aligned} P_2(b) &= -\int_a^b \left( \int_a^t G_1(x)(x-t)^{n-2} dx \right) dt \\ &= \int_a^{a+\lambda} \left( \int_a^x (1-g(s)) ds \right) \frac{(x-b)^{n-1}}{n-1} dx + \int_{a+\lambda}^b \left( \int_x^b g(s) ds \right) \frac{(x-b)^{n-1}}{n-1} dx \\ &= \int_a^{a+\lambda} (x-a) \frac{(x-b)^{n-1}}{n-1} dx + \lambda \cdot \int_{a+\lambda}^b \frac{(x-b)^{n-1}}{n-1} dx - \int_a^b \left( \int_a^x g(s) ds \right) \frac{(x-b)^{n-1}}{n-1} dx \\ &= \frac{(a-b)^{n+1}}{(n-1) \cdot n \cdot (n+1)} - \frac{(a+\lambda-b)^{n+1}}{(n-1) \cdot n \cdot (n+1)} + \int_a^b g(x) \frac{(x-b)^n}{(n-1) \cdot n} dx \end{aligned}$$

and

$$\begin{aligned} m_2 &= -\frac{1}{P_2(b)} \int_a^b \left( \int_a^t G_1(x)(x-t)^{n-2} dx \right) t dt \\ &= -\frac{1}{P_2(b)} \int_a^b G_1(x) \left( \int_x^b (x-t)^{n-2} \cdot t dt \right) dx \\ &= -\frac{1}{P_2(b)} \int_a^b G_1(x) \left( t \cdot \frac{-(x-t)^{n-1}}{n-1} \Big|_x^b + \int_x^b \frac{(x-t)^{n-1}}{n-1} dt \right) dx \\ &= -\frac{1}{P_2(b)} \int_a^b G_1(x) \left( -b \cdot \frac{(x-b)^{n-1}}{n-1} - \frac{(x-b)^n}{(n-1) \cdot n} \right) dx \\ &= b + \frac{1}{P_2(b)} \int_a^b G_1(x) \frac{(x-b)^n}{(n-1) \cdot n} dx \\ &= b + \frac{1}{(n-1) \cdot n \cdot (n+1) \cdot P_1(b)} \\ &\quad \times \left( \frac{(a-b)^{n+2}}{n+2} - \frac{(a+\lambda-b)^{n+2}}{n+2} + \int_a^b g(x)(x-b)^{n+1} dx \right). \end{aligned}$$

Hence, the proof is completed.  $\square$

**Theorem 7.** Let  $f: [a, b] \rightarrow \mathbb{R}$  be  $(n + 2)$ -convex on  $[a, b]$  and  $f^{(n-1)}$  absolutely continuous for  $n \geq 2$ . Let  $g: [a, b] \rightarrow \mathbb{R}$  be an integrable function such that  $0 \leq g \leq 1$  and  $\lambda = \int_a^b g(t)dt$ . Let function  $G_2$  be defined by the following.

$$G_2(x) = \begin{cases} \int_a^x g(t)dt, & x \in [a, b - \lambda], \\ \int_x^b (1 - g(t))dt, & x \in [b - \lambda, b]. \end{cases} \tag{13}$$

Then, the following is obtained:

$$\begin{aligned} &P_3(b) \cdot f^{(n)}(m_3) \leq \\ &(n - 2)! \left[ \int_{b-\lambda}^b f(t)dt - \sum_{i=0}^{n-2} \frac{f^{(i+1)}(a)}{i!} \int_a^b G_2(x)(x - a)^i dx - \int_a^b f(t)g(t)dt \right] \\ &\leq P_3(b) \cdot \left[ \frac{b - m_3}{b - a} f^{(n)}(a) + \frac{m_3 - a}{b - a} f^{(n)}(b) \right], \end{aligned} \tag{14}$$

where

$$P_3(b) = \frac{1}{(n - 1) \cdot n} \left( \frac{(b - a)^{n+1} - (b - \lambda - a)^{n+1}}{n + 1} - \int_a^b g(x)(x - a)^n dx \right)$$

and

$$\begin{aligned} m_3 = a + &\frac{1}{(n - 1) \cdot n \cdot (n + 1) \cdot P_3(b)} \\ &\times \left( \frac{(b - a)^{n+2} - (b - \lambda - a)^{n+2}}{n + 2} - \int_a^b g(x)(x - a)^{n+1} dx \right). \end{aligned}$$

**Proof.** We follow the similar arguments as in the proof of Theorem 5. As function  $f^{(n-1)}$  is absolutely continuous, the identity (5) holds. The inequality (14) follows directly from Theorem 1, substituting the non-negative function  $p$  by a non-negative function of the following:

$$p(t) = \int_t^b G_2(x)(x - t)^{n-2} dx$$

and a convex function  $f$  by a convex function  $f^{(n)}$ , and then using identity (5) for integral  $\int_a^b \left( \int_t^b G_2(x)(x - t)^{n-2} dx \right) f^{(n)}(t)dt$ . Furthermore, we calculate  $P_3(b)$  and  $m_3$  as follows.

$$\begin{aligned} P_3(b) &= \int_a^b \left( \int_t^b G_2(x)(x - t)^{n-2} dx \right) dt \\ &= \int_a^{b-\lambda} \left( \int_a^x g(s)ds \right) \frac{(x - a)^{n-1}}{n - 1} dx + \int_{b-\lambda}^b \left( \int_x^b (1 - g(s))ds \right) \frac{(x - a)^{n-1}}{n - 1} dx \\ &= \int_{b-\lambda}^b (b - x) \frac{(x - a)^{n-1}}{n - 1} dx - \lambda \cdot \int_{b-\lambda}^b \frac{(x - a)^{n-1}}{n - 1} dx + \int_a^b \left( \int_a^x g(s)ds \right) \frac{(x - a)^{n-1}}{n - 1} dx \\ &= \frac{(b - a)^{n+1} - (b - \lambda - a)^{n+1}}{(n - 1) \cdot n \cdot (n + 1)} - \int_a^b g(x) \frac{(x - a)^n}{(n - 1) \cdot n} dx, \end{aligned}$$



$$\begin{aligned}
 m_3 &= \frac{1}{P_3(b)} \int_a^b \left( \int_t^b G_2(x)(x-t)^{n-2} dx \right) t dt \\
 &= \frac{1}{P_3(b)} \int_a^b G_2(x) \left( \int_a^x (x-t)^{n-2} \cdot t dt \right) dx \\
 &= \frac{1}{P_3(b)} \int_a^b G_2(x) \left( t \cdot \frac{-(x-t)^{n-1}}{n-1} \Big|_a^x + \int_a^x \frac{(x-t)^{n-1}}{n-1} dt \right) dx \\
 &= \frac{1}{P_3(b)} \int_a^b G_2(x) \left( \frac{a \cdot (x-a)^{n-1}}{n-1} + \frac{(x-a)^n}{(n-1) \cdot n} \right) dx \\
 &= a + \frac{1}{P_3(b)} \int_a^b G_2(x) \frac{(x-a)^n}{(n-1) \cdot n} dx \\
 &= a + \frac{1}{P_3(b)} \left( \frac{(b-a)^{n+2} - (b-\lambda-a)^{n+2}}{(n-1) \cdot n \cdot (n+1) \cdot (n+2)} - \int_a^b g(x) \frac{(x-a)^{n+1}}{(n-1) \cdot n \cdot (n+1)} dx \right).
 \end{aligned}$$

Hence, the proof is completed.  $\square$

**Theorem 8.** Let  $f: [a, b] \rightarrow \mathbb{R}$  be  $(n + 2)$ -convex on  $[a, b]$  and  $f^{(n-1)}$  absolutely continuous for  $n \geq 2$ . Let  $g: [a, b] \rightarrow \mathbb{R}$  be an integrable function such that  $0 \leq g \leq 1$  and  $\lambda = \int_a^b g(t)dt$ . Let function  $G_2$  be defined by (13). If the following is the case:

$$\int_a^t G_2(x)(x-t)^{n-2} dx \leq 0, \quad t \in [a, b]$$

then we obtain the following:

$$\begin{aligned}
 &P_4(b) \cdot f^{(n)}(m_4) \leq \\
 &(n-2)! \left[ \int_{b-\lambda}^b f(t)dt - \sum_{i=0}^{n-2} \frac{f^{(i+1)}(b)}{i!} \int_a^b G_2(x)(x-b)^i dx - \int_a^b f(t)g(t)dt \right] \tag{15} \\
 &\leq P_4(b) \cdot \left[ \frac{b-m_4}{b-a} f^{(n)}(a) + \frac{m_4-a}{b-a} f^{(n)}(b) \right],
 \end{aligned}$$

where

$$P_4(b) = \frac{-1}{(n-1) \cdot n} \left( \frac{(-\lambda)^{n+1}}{n+1} + \int_a^b g(x)(x-b)^n dx \right)$$

and

$$m_4 = b - \frac{1}{(n-1) \cdot n \cdot (n+1) \cdot P_4(b)} \left( \frac{(-\lambda)^{n+2}}{n+2} + \int_a^b g(x)(x-b)^{n+1} dx \right).$$

**Proof.** Under the assumption that  $\int_a^t G_2(x)(x-t)^{n-2} dx \leq 0$ , it is obvious that the following is the case:

$$p(t) = - \int_a^t G_2(x)(x-t)^{n-2} dx \tag{16}$$

where it is a non-negative function. Again, replacing  $p(t)$  in Theorem 1 by (16) and  $f$  by  $f^{(n)}$  and then using the identity (6) for

$$\int_a^b \left( \int_a^t G_2(x)(x-t)^{n-2} dx \right) f^{(n)}(t)dt,$$

we obtain the required inequalities (15). Finally, a simple calculation yields the following:

$$\begin{aligned}
 P_4(b) &= - \int_a^b \left( \int_a^t G_2(x)(x-t)^{n-2} dx \right) dt \\
 &= \int_a^{b-\lambda} \left( \int_a^x g(s) ds \right) \frac{(x-b)^{n-1}}{n-1} dx + \int_{b-\lambda}^b \left( \int_x^b (1-g(s)) ds \right) \frac{(x-b)^{n-1}}{n-1} dx \\
 &= - \int_{b-\lambda}^b \frac{(x-b)^n}{n-1} dx - \lambda \cdot \int_{b-\lambda}^b \frac{(x-b)^{n-1}}{n-1} dx + \int_a^b \left( \int_a^x g(s) ds \right) \frac{(x-b)^{n-1}}{n-1} dx \\
 &= - \frac{(-\lambda)^{n+1}}{(n-1) \cdot n \cdot (n+1)} - \int_a^b g(x) \frac{(x-b)^n}{(n-1) \cdot n} dx
 \end{aligned}$$

and

$$\begin{aligned}
 m_4 &= \frac{-1}{P_4(b)} \int_a^b \left( \int_a^t G_2(x)(x-t)^{n-2} dx \right) t dt \\
 &= \frac{-1}{P_4(b)} \int_a^b G_2(x) \left( \int_x^b (x-t)^{n-2} \cdot t dt \right) dx \\
 &= \frac{-1}{P_4(b)} \int_a^b G_2(x) \left( t \cdot \frac{-(x-t)^{n-1}}{n-1} \Big|_x^b + \int_x^b \frac{(x-t)^{n-1}}{n-1} dt \right) dx \\
 &= \frac{-1}{P_4(b)} \int_a^b G_2(x) \left( -b \cdot \frac{(x-b)^{n-1}}{n-1} - \frac{(x-b)^n}{(n-1) \cdot n} \right) dx \\
 &= b + \frac{1}{P_4(b)} \int_a^b G_2(x) \frac{(x-b)^n}{(n-1) \cdot n} dx \\
 &= b - \frac{1}{P_4(b)} \left( \frac{(-\lambda)^{n+2}}{(n-1) \cdot n \cdot (n+1) \cdot (n+2)} + \int_a^b g(x) \frac{(x-b)^{n+1}}{(n-1) \cdot n \cdot (n+1)} dx \right).
 \end{aligned}$$

□

**Remark 1.** If function  $f$  is  $(n + 2)$ -concave, the inequalities in Theorems 5–8 are reversed. This follows from the fact that for  $(n + 2)$ -concave function, we have  $-f^{(n+2)} \geq 0$ . Hence,  $-f^{(n)}$  is convex and we can apply inequality (1) to function  $-f^{(n)}$ .

**Remark 2.** The expressions  $P_i(b)$  and  $m_i$  for  $i = 1, \dots, 4$  can also be achieved by the method introduced in [16]. By this method, we calculate  $P_1(b)$  and  $m_1$ . Other expressions can be recaptured in a similar manner.

The value of  $P_1(b)$  can be obtained from (3) by taking  $f(t) = \frac{(t-a)^n}{n!}$ . Then,  $f^{(n)}(t) = 1$ . Thus, we have the following.

$$\begin{aligned}
 P_1(b) &= -(n-2)! \left( \int_a^{a+\lambda} \frac{(x-a)^n}{n!} dt - \int_a^b \frac{(x-a)^n}{n!} g(t) dt \right) \\
 &= - \frac{\lambda^{n+1}}{(n-1) \cdot n \cdot (n+1)} + \int_a^b \frac{(x-a)^n}{(n-1) \cdot n} g(t) dt.
 \end{aligned}$$

Hence, we obtained expression (9).

From Theorem 1, we previously obtained the following.

$$m_1 = \frac{1}{P_1(b)} \int_a^b \left( \int_t^b G_1(x)(x-t)^{n-2} dx \right) t dt.$$

To calculate  $m_1$ , we take function  $f(t) = \frac{(t-a)^{n+1}}{(n+1)!}$ . Then,  $f^{(n)}(t) = t - a$ . Hence, from the identity (3), we obtain expression (10).

### 3. Conclusions

In this paper, we obtained new weighted Hermite–Hadamard-type inequalities for higher order convex functions. We used previously obtained identities related to the generalizations of Steffensen’s inequality. Results obtained in this paper can be considered as a starting point for some future work.

**Author Contributions:** Conceptualization, J.P., A.P.P. and K.S.K.; Writing — original draft, J.P., A.P.P. and K.S.K. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** Croatian Science Foundation (HRZZ 7926) “Separation of parameter influence in engineering modeling and parameter identification”.

**Conflicts of Interest:** The authors declare no conflict of interest.

### References

1. Hermite, C. Sur deux limites d’une intégrale dé finie. *Mathesis* **1883**, *3*, 82.
2. Hadamard, J. Étude sur les propriétés des fonctions entières et en particulier d’une fonction considérée par Riemann. *J. Math. Pures Appl.* **1893**, *58*, 171–215.
3. Dinu, C. A weighted Hermite Hadamard inequality for Steffensen-Popoviciu and Hermite- Hadamard weights on time scales, *Analele Stiintifice ale Universitatii Ovidius Constanta–Seria. Matematica* **2009**, *17*, 77–90.
4. Abbas, M.I.; Ragusa, M.A. Solvability of Langevin equations with two Hadamard fractional derivatives via Mittag-Leffler functions. *Appl. Anal.* **2021**, 1–15. [\[CrossRef\]](#)
5. Niculescu, C.P.; Persson, L.-E. Old and new on the Hermite-Hadamard inequality. *Real Anal. Exch.* **2004**, *29*, 663–685. [\[CrossRef\]](#)
6. Niculescu, C.P.; Persson, L.-E. *Convex Functions and Their Applications: A Contemporary Approach*; CMS Books in Mathematics; Springer: New York, NY, USA, 2005.
7. Niculescu, C.P.; Stanescu, M.M. The Steffensen-Popoviciu measures in the context of qualiconvex functions. *J. Math. Inequal.* **2017**, *11*, 469–483. [\[CrossRef\]](#)
8. Pečarić, J.E.; Proschan, F.; Tong, Y.L. Convex functions, partial orderings, and statistical applications. In *Mathematics in Science and Engineering 187*; Academic Press: San Diego, CA, USA, 1992.
9. Pečarić, J.; Perić, I. Refinements of the integral form of Jensen’s and Lah-Ribarić inequalities and applications for Csiszár divergence. *J. Inequal. Appl.* **2020**, *108*, 287 [\[CrossRef\]](#)
10. Wu, S. On the Weighted Generalization of the Hermite-Hadamard Inequality and Its Applications. *Rocky Mountain J. Math.* **2009**, *39*, 1741–1749. [\[CrossRef\]](#)
11. Steffensen, J.F. On certain inequalities between mean values and their application to actuarial problems. *Skand. Actuar. J.* **1918**, *1918*, 82–97. [\[CrossRef\]](#)
12. Jakšetić, J.; Pečarić, J.; Perušić Pribanić, A.; Smoljak Kalamir, K. *Weighted Steffensen’s Inequality (Recent Advances in Generalizations of Steffensen’s Inequality)*; Monographs Inequalities 17: Element, Zagreb, 2020.
13. Pečarić, J.; Smoljak Kalamir, K.; Varošaneć, S. Steffensen’s and Related Inequalities (A Comprehensive Survey and Recent Advances). *Monographs in Inequalities 7*. 2014. Available online: <http://ele-math.com/static/pdf/books/593-mia07.pdf> (accessed on 29 March 2022).
14. Mitrinović, D.S. *The Steffensen Inequality*; Publikacije Elektrotehničkog Fakulteta. Serija Matematika i Fizika No. 247/273; University of Belgrade: Belgrade, Serbia, 1969; pp. 1–14.
15. Pečarić, J.; Perušić Pribanić, A.; Smoljak Kalamir, K. Generalizations of Steffensen’s inequality via Taylor’s formula. *J. Inequal. Appl.* **2015**, *207*, 1–25. [\[CrossRef\]](#)
16. Barić, J.; Kvesić, L.J.; Pečarić, J.; Ribičić Penava, M. Fejér type inequalities for higher order convex functions and quadrature formulae. *Aequat. Math.* **2021**, *96*, 417–430. [\[CrossRef\]](#)